
Phago-ENKO Bacteriophage

Product description

Composition:

Phago-Enko bacteriophage is a sterile filtrate of phage lysates active to:

- Shigella;
- Salmonella;
- Different types of E. coli;
- Species of pathogenic Staphylococcus;

Inactive ingredients: Bacteriological growth medium, standard saline solution. Chinazolin as a preservative.

Description:

Preparation is transparent liquid, with yellow to light brown colour.

Pharmacological actions:

Phago-Enko Bacteriophage leads to a specific lysis of bacteria belonging to Staphylococcus, Shigella, E.coli and Salmonella of different serovars.

Recommendations for use:

Phago-ENKO bacteriophage is a paratherapeutic product. It can be used during enteric infections caused by the above-mentioned microorganisms as well as for prophylactic purposes.

Indications:

Enteric infections: dysentery, salmonellosis, dyspepsia, colitis, gastro enterocolitis, cholecystitis, staphylococcal toxic infection, disbiosis etc.

Usage and dosage:

Phago-Enko phage can be used per os from the first days of the disease for 5 - 6 days in the following doses:

- Children under 3 years - 5ml per reception,
- Children above 3 years old and adolescents - 10 ml at once, 4 times a day, 30 minutes before the meal.

Before taking the phage per os, taking 2-3% solution of baking soda (one glass, 150-200 ml) is recommended to reduce stomach acid.

During infectious intestinal diseases, the solution is given in combination with applying the phage by enemas; preferably, in the evening time once a day, after a cleansing enema with a baking soda solution, per rectum: 10 ml of the preparation for children and 20 ml - for adults.

During outbreak of infectious intestinal diseases, the solution is given to all patients, regardless of clinical diagnosis, 10-20 ml twice a day, for five days.

Special Warnings:

Before use, the vial should be well shaken. The contents should be transparent liquid without sediment.

The solution should not be used in case of vials damage or if the liquid inside is non-transparent!

Do not store and use the solution after destroying the integrity of the vial!!!

Overdosing, contradictions and side effects:

During the usage of Phago-ENKO bacteriophage, overdosing symptoms have not been revealed. Usage of Phago-ENKO bacteriophage does not have contradictions. No side effects have been revealed during application of Phago-ENKO bacteriophage.

Usage together with other preparations:

Phago-ENKO can be used in parallel or supplemental to other medications. Drinking alcohol during use of bacteriophages is not recommended.

Specific categories of patients:**Pregnancy and lactation:**

Bacteriophages can be used during pregnancy and lactation period.

Usage in elderly people:

Bacteriophages can be used in elderly people.

Effect on drivers and people working with potentially harmful machines:

Usage of bacteriophages does not affect driving skills or skills of other equipment usage.

Packaging:

Phago-ENKO bacteriophage is produced in 10 ml vials (5 vials per pack).

Storage conditions and shelf life:

Store at +2°C - +8°C in dry place, protected from direct light.

Shelf life 24 month.

Issued without prescription

Phago-ENKO is a paratherapeutic product and doesn't include GMO.

Manufactured by:

"Eliava BioPreparations" LLC

3, Gotua str. Tbilisi, 0160, Georgia

Phone: +995 (32) 2151402

E-mail: info@phage.ge