

Phago-Intesti Bacteriophage

Product description

Composition:

Intesti bacteriophage is a mix of sterile filtrate of phage lysates active against:

- Shigella;
- Salmonella;
- Different types of E. coli;
- Proteus;
- Staphylococcus;
- Pseudomonas aeruginosa;
- Enterococcus;

Inactive ingredients: Bacteriological growth medium, standard sodium saline. Chinazolin as preservative.

Description:

Solution is a transparent liquid, with yellow to light brown colour.

Pharmacological actions:

Phago-Intesti bacteriophage leads to specific lysis of following bacteria: Shigella; Salmonella; Escherichia coli; Proteus; Staphylococcus; Pseudomonas aeruginosa; Enterococcus.

Recommendations for use:

Phago-Intesti bacteriophage is a parapharmaceutical product. It can be used during intestinal infectious diseases caused by enlisted microorganisms and for the prophylaxis.

Indications:

- Dysentery;
- Salmonellosis;
- Dyspepsia.;
- Infective colitis;
- Enterocolitis;
- Dysbiosis;

Usage and dosage:

Phago-Intesti bacteriophage can be used per os with first signs of disease within 5-6 days. Phago-Intesti bacteriophage should be used in following doses:

Age	Amount per admission	Period of admissions
Children up to 3 years of age	5 ml per admission	4 times a day 30 minutes before meal
Children from age of 3 and adults	10 ml per admission	4 times a day 30 minutes before meal

During Per os application of Phago- Intesti bacteriophage in adults and children from age of 3, drinking of 2-3% baking soda solution (one glass, 150-250 ml) is recommended.

In parallel to per os admission of Phago-Intesti bacteriophage it is recommended to use it per rectum, once a day in the amount of 10 ml for children and 20 ml for adults. (Per rectum usage is recommended in the evening with enema after emptying stomach)

For prophylaxis use Phago-Intesti bacteriophage is recommended:

- In organized groups to prevent spread of infections. During seasonal infections, Phago-Intesti bacteriophage can be applied once a day in amount of 10-20 ml. Application can be made with the interval

Revision 1

Revision Date: 10.04.2024

of 5 days during 1 month.

- During the increase cases of intestinal infections, all patients in the clinics are recommended to go through phage treatment despite diagnoses. Phage application is recommended 2 times with the interval of 5 days. Amount per admission should be 10-20 ml.

Special Warnings:

Before use, the vial should be well shaken. The contents should be transparent liquid without sediment. The solution should not be used in case of vials damage or if the liquid inside is non-transparent! Do not store and use the solution after destroying the integrity of the vial!!!

Overdosing, contradictions and side effects:

During the usage of Phago-Intesti bacteriophage, overdosing symptoms have not been revealed. Usage of Phago-Intesti bacteriophage does not have contradictions. No side effects have been revealed during usage of Phago-Intesti bacteriophage.

Usage together with other preparations:

Phago-Intesti can be used in parallel or supplemental to other medications. Drinking alcohol during use of bacteriophages is not recommended.

Specific categories of patients:

Pregnancy and lactation:

Phago-Intesti can be used during pregnancy and lactation period.

Usage in elderly people:

Phago-Intesti is used in elderly people.

Effect on drivers and people working with potentially harmful machines:

Usage of bacteriophages does not affect driving skills or skills of other equipment usage.

Packaging:

Phago-Intesti bacteriophage is produced in 10 ml vials (5 vials per pack).

Storage conditions and shelf life:

Store at +2°C - +8°C in dry place, protected from direct light.
Shelf life 24 month.

Issued without prescription

Developed by G. Eliava Institute of Bacteriophages, Microbiology and Virology.

Phago-Intesti is a parapharmaceutical product and doesn't include GMO.

Manufactured by:

"Eliava BioPreparations" LLC
3, Gotua str. Tbilisi, 0160, Georgia
Phone: +995 (32) 2370702
E-mail: info@phage.ge